Extra generations and discrepancies of electroweak precision data

V.A. Novikov, L.B. Okun,
ITEP, Moscow, Russia
A.N. Rozanov,
CPPM, IN2P3, CNRS, Univ. Mediteranee, Marseilles, France
and ITEP, Moscow, Russia
M.I. Vysotsky
ITEP, Moscow, Russia

Abstract

It is shown that additional chiral generations are not excluded by the latest electroweak precision data if one assumes that there is no mixing with other three generations. In the case of "heavy extra generations", when all four new particles are heavier than Z-boson, quality of the fit for the one new generation is as good as for zero new generations (Standard Model). In the case of neutral leptons with masses around 50 GeV ("partially heavy extra generation") the minimum of χ^2 is between one and two extra generations.

Two years ago in paper [1] we analyzed bounds on non-decoupled New Physics in a form of additional heavy quark-lepton generations from electroweak precision data. It was shown that while the case of all four new fermions (U and D quarks, neutral lepton N and charged lepton E) heavier than Z-boson was excluded at 2.5 σ level, existence of new generations with relativity light neutral lepton N ($m_N \approx 50$ GeV) was allowed. At that time quality of Standard Model (SM) fit of the data was very good, $\chi^2/n_{\rm d.o.f.} = 15/14$. At the time of Osaka Conference, summer 2000, nothing radical happened but χ^2 became 21/14 and the level at which one extra heavy generation was excluded went down to 2σ [2]. However the latest precision data announced summer 2001 [3] has changed the situation: the fit is still bad, 24/13, but now the presence of one heavy generation does not make the fit worse as compared with SM.

In Table 1 LEPTOP fit of summer 2001 data is presented. There are two most significant changes in comparison with previous data presented in Table 2:

1. Due to precision measurement of the cross-section of e^+e^- annihilation into hadrons in the interval 2-5 GeV at BES the error in $\bar{\alpha} \equiv \alpha(M_Z)$ is now two times smaller. (Following Electroweak Working Group (EWWG) we use result [4] though other estimates can be found in the literature as well);

Table 1: LEPTOP fit to electroweak observables. Year 2001. By italics we designate calculated (not measured) quantities.

Source	Observable	Exp. data	LEPTOP fit	Pull
LEP I	Γ_Z [GeV]	2.4952(23)	2.4966(16)	-0.6
	σ_h [nb]	41.540(37)	41.480(14)	1.6
	A_{FB}^l	0.0171(10)	0.0165(3)	0.7
	R_l	20.767(25)	20.738(18)	1.1
	A_τ, A_e	0.1465(33)	0.1483(11)	-0.5
	R_b	0.2165(7)	0.2157(1)	1.2
	R_c	0.1719(31)	0.1723(1)	-0.1
	A_{FB}^b	0.0990(17)	0.1040(8)	-2.9
	A_{FB}^c	0.0685(34)	0.0743(6)	-1.7
	$s_l^2(Q_{FB})$	0.2324(12)	0.2314(1)	0.9
SLC	A_{LR}	0.1513(21)	0.1483(11)	1.4
	s_l^2 (A_{LR})	*0.2310(3)*	*0.2314(1)*	-1.4
	A_b	0.9220(200)	0.9349(1)	-0.6
	A_c	0.6700(260)	0.6684(5)	0.1
LEP II, Tevatron	m_W [GeV]	80.451(33)	80.392(20)	1.8
	$s_W^2(m_W)$	*0.2216(6)*		
Tevatron	s_W^2 (νN)	0.2255(21)	0.2230(3)	1.2
	$m_W(\nu N)$ [GeV]	*80.250(109)*		
	m_t [GeV]	174.3(5.1)	175.0(4.4)	-0.1
Fit	m_H [GeV]		79^{+47}_{-29}	
	$\hat{\alpha}_s$		0.1182(27)	
$e^+e^- \to$ hadrons	$\bar{\alpha}^{-1}$	128.936(49)	128.918(45)	0.4
	$\chi^2/n_{d.o.f.}$		23.8/13	

2. Central value of M_W is now bigger by a half of σ.

The latter is the main cause for the relaxation of the bound on heavy extra generations.

Exclusion plot for the number N_g of extra heavy generations is presented in Fig. 1.

To produce this plot we take $m_D = 130$ GeV – the Tevatron lower bound on new quark mass; we use experimental 95% C.L. bound on higgs mass $m_H > 113$ GeV [3] and vary $\Delta m = \sqrt{m_U^2 - m_D^2}$ and number of extra generations N_g. (In order to have two-dimensional plot we arbitrary assumed that $m_N = m_U$ and $m_E = m_D$; other choices do not change the obtained results drastically); χ^2 minimum corresponds to unphysical point $N_g = 0.5$. For 170 GeV $< m_U <$ 200 GeV we get the same quality of fit in the case $N_g = 1$ as that for the Standard Model ($N_g = 0$). In ref. [5] one can find a statement that extra heavy generations are excluded by the precision electroweak data. However, analysis

Table 2: LEPTOP fit to electroweak observables. Year 2000.

Source	Observable	Exp. data	LEPTOP fit	Pull
LEP I	Γ_Z [GeV]	2.4952(23)	2.4964(16)	-0.5
	σ_h [nb]	41.541(37)	41.479(15)	1.7
	A_{FB}^l	0.0171(10)	0.0164(3)	0.7
	R_l	20.767(25)	20.739(18)	1.1
	A_τ, A_e	0.1467(32)	0.1480(13)	-0.4
	R_b	0.2165(7)	0.2157(1)	1.2
	R_c	0.1709(34)	0.1723(1)	-0.4
	A_{FB}^b	0.0990(20)	0.1038(9)	-2.4
	A_{FB}^c	0.0689(35)	0.0742(7)	-1.5
	$s_l^2(Q_{FB})$	0.2321(10)	0.2314(2)	0.7
SLC	A_{LR}	0.1514(22)	0.1480(16)	1.5
	s_l^2 (A_{LR})	*0.2310(3)*	*0.2314(2)*	-1.5
	A_b	0.9110(250)	0.9349(1)	-1.0
	A_c	0.6300(260)	0.6683(6)	-1.5
LEP II, Tevatron	m_W [GeV]	80.434(37)	80.397(23)	1.0
	$s_W^2(m_W)$	*0.2219(7)*		
Tevatron	s_W^2 (νN)	0.2255(21)	0.2231(2)	1.1
	$m_W(\nu N)$ [GeV]	*80.250(109)*		
	m_t [GeV]	174.3(5.1)	174.0(4.2)	0.1
Fit	m_H [GeV]		55^{+45}_{-26}	
	$\hat{\alpha}_s$		0.1183(27)	
$e^+e^- \to$ hadrons	$\bar{\alpha}^{-1}$	128.878(90)	128.850(90)	0.3
	$\chi^2/n_{d.o.f.}$		21.4/14	

performed in [5] refers to upper and lower parts of Fig. 1, $\Delta m > 200$ GeV and $\Delta m = 0$, where existence of new heavy generations is really strongly suppressed. This is not the case for the central part of Fig. 1 ($\Delta m \approx 150$ GeV).

Two heavy generations are excluded at more than 3σ level. Nevertheless, two and even three "partially heavy" generations are allowed when neutral fermions are relatively light, $m_N \simeq 55$ GeV (see Fig. 2). Using all existing LEP II statistics on the reactions $e^+e^- \to \gamma + \nu\bar{\nu}, \gamma + N\bar{N}$ in dedicated search one can exclude 3 "partially heavy" generations which contain such a light N at a level of 3σ (see [6]), while one or even two such generations may exist.

The cause of disappearance of the suppression of extra heavy generations which existed in the early data is the contradiction in description of modern data on M_W and s_l^2 in the framework of SM. The point is that the only free parameter of SM, higgs boson mass, has the following values being extracted from these observables:

$$\begin{aligned} (m_W)_{\rm LEPII,Tevatron,NuTeV} &= 80.428(32)\ {\rm GeV} \Rightarrow \\ &\Rightarrow \quad m_H = 50^{+50}_{-35}\ {\rm GeV}\ , \end{aligned} \tag{1}$$



Figure 1: Exclusion plot for heavy extra generations. χ^2 minimum is shown by cross; borders of regions show domains allowed at the level $1\sigma, 2\sigma$, etc. N_g is the number of extra generations.

$$\begin{aligned} (s_l^2)_{\rm LEPI,SLAC} &= 0.23140(15)\ {\rm GeV} \Rightarrow \\ &\Rightarrow \quad m_H = 150^{+75}_{-50}\ {\rm GeV}\ . \end{aligned} \tag{2}$$

$N_g = 0.5$ reduces the contradiction between the two values of m_H. Nevertheless the resulting χ^2 does not improve drastically and this is due to another "defect" of precision data: the discrepancy between the average value of s_l^2 extracted from pure leptonic measurements and its value from events with hadrons in final state [3]:

	s_l^2
Leptons	0.23113(21)
Hadrons	0.23230(29)

(3)

These 3.3 σ difference is the root of poor quality of the SM fit. The hadronic measurements are dominated by very small uncertainty of the forward-backward asymmetry in reaction $e^+e^- \to Z \to b\bar{b}$. According to Table 1

$$(A_{FB}^b)_{\rm exp} = 0.0990(17)\ . \tag{4}$$

One can question whether such a good accuracy can be obtained in the analysis of hadronic jets production. Another value of A_{FB}^b can be obtained by multiplying measured at SLAC beauty asymmetry A_b and leptonic asymmetry



Figure 2: Exclusion plot for the number of partially heavy extra generations with light neutral lepton N. On horizontal axis – N_g, on vertical axis – the mass of the neutral lepton m_N.

A_l. Then

$$A_{FB}^b = \frac{3}{4} A_b A_l = 0.1038(25) \ . \tag{5}$$

The number (5) differs from (4), but nicely coincides with the SM fit: 0.1040(8) (see Table 1).

Let us assume following Chanowitz [7] that A_{FB}^b has larger uncertainty than given in Eq. (4) and look to what consequences in respect of extra generations this hypothesis will lead.[1] If we multiply experimental uncertainties of A_{FB}^b and A_{FB}^c, which are strongly correlated, by a factor 10, the quality of SM fit improves drastically: $\chi^2/n_{d.o.f.}$ shifts from 23.8/13 to 10.9/13 and simultaneously one heavy extra generation becomes excluded at the level of 2.5σ (see Fig. 3).

However, a serious problem arises: it is just A_{FB}^b given by Eq. (4) which pushes m_H to larger values. With our modification of experimental results on A_{FB}^b and A_{FB}^c the SM fit gives:

$$m_H = 42^{+30}_{-18} \text{ GeV} \ , \tag{6}$$

[1] Another way to resolve situation with A_{FB}^b is to assume that there exist New Physics contributions to $Zb\bar{b}$ couplings g_V^b and g_A^b. Since in the expression for A_{FB}^b the corrections to g_V^b and g_A^b are multiplied by small factor g_V^e they should be large, so they must appear at the tree level. Also $Z \to b\bar{b}$ width proportional to $(g_A^b)^2 + (g_V^b)^2$ should not noticeably change since $R_b \equiv \Gamma_{Z\to b\bar{b}}/\Gamma_Z$ is at present in good agreement with SM fit, see Table 1. In recent paper [8] inclusion of additional bottom-like heavy quarks with vector currents is suggested to resolve the discrepancy (3).



Figure 3: Exclusion plot for heavy extra generations with 10 times enlarged errors in A_{FB}^b and A_{FB}^c. χ^2 minimum is at the upper border of the Fig.

well below modern LEP II bound: $m_H > 113$ GeV, a substantial trouble for the SM.

What concerns partially heavy extra generations, they nicely fit data even with ten times enlarged uncertainties of A_{FB}^b and A_{FB}^c, see Fig. 4. At both minima $\chi^2/n_{d.o.f.} \approx 13/13$, while $m_H \approx 40$ GeV.

In the recent paper [9] it was noted that SUSY extension of Standard Model with light sneutrinos with masses in the range 55- 80 GeV is allowed by precision data and pushes higgs mass to larger values.

The presence of new heavy particles is important for production and decay of higgs boson. New quarks considerably enhance higgs production at Tevatron and LHC through gluon fusion which should help to discover this particle [10]. If higgs decay into a pair of neutral leptons is kinematically allowed it will dominate, so that a moderately heavy higgs will decay invisibly [11]. At LEP II the invisibly decaying higgs is excluded at the same level as the SM higgs by missing mass method [12]. Contrary to that the LHC will look for visible decay modes of higgs. If the branching ratios of the latter are small this will be not easy.

We are grateful to A. Olshevsky for providing us with ref. [3] and to P.S. Bambade and M. Stanitzki for providing us with ref. [12].

L.O., A.R. and M.V. are grateful to CERN EP and TH for hospitality; L.O., V.N. and M.V. were partly supported by RFBR grant No. 00-15-96562.



Figure 4: Exclusion plot for partially heavy extra generations with 10 times enlarged errors in A_{FB}^b and A_{FB}^c. Two local χ^2 minima are shown.

References

[1] M. Maltoni, V.A. Novikov, L.B. Okun, A.N. Rozanov, and M.I. Vysotsky, Phys. Lett. **B476** (2000) 107.

[2] V.A. Ilyin, M. Maltoni, V.A. Novikov, L.B. Okun, A.N. Rozanov, and M.I. Vysotsky, *in:* Proceedings of the XXX International Conference on High Energy Physics, Osaka-2000, World Scientific, v.2, p. 1209-1212.

[3] M.W. Grünewald, Talk at LEP Physics Jamboree, CERN, July 10th, 2001; http://www.cern.ch/LEPEWWG.

[4] H. Burkhardt and B. Pietrzyk, LAPP-EXP-2001-03.

[5] J. Erler and P. Langacker, *in:* Review of Particle Physics, The European Physical Journal **15** (2000) 95.

[6] V.A. Ilyin et al., Phys. Lett. **B503** (2001) 126.

[7] M.S. Chanowitz, hep-ph/0104024 (2001).

[8] D. Choudhury, T.M.P. Tait and C.E.M. Wagner, hep-ph/0109097 (2001).

[9] G. Altarelli, F. Caravaglios, G.F. Giudice, P. Gambino, G. Ridolfi, JHEP 0106 (2001) 018.

[10] V.D. Barger and R.J.N. Philips, *Collider Physics*, Addison-Wesley (1997);
E. Arik et al., CERN-ATLAS Internal Note, ATL-PHUS-98-125 (1998);
I.F. Ginzburg, I.P. Ivanov and A. Schiller, Phys. Rev. **D60** (1999) 095001;
E. Arik et al., hep-ph/0109037 (2001).

[11] D. Fargion, Yu.A. Golubkov, M.Yu. Khlopov, R.V. Konoplich, R. Migrani,
JETP Lett. **69** (1999) 434; Pis'ma v ZhETP **69** (1999) 402;
V.A. Khoze, hep-ph/0105069.

[12] ALEPH, DELPHI, L3 and OPAL Collaborations, The LEP working group for Higgs boson searches; ALEPH 2001-36, DELPHI 2001-117, L3 note 2702, Opal Technical Note TN694; paper submitted to EPS01 in Budapest.